

Mass Shifts through Re-scattering

Ron S. Longacre

Brookhaven National Laboratory, Upton, NY 11973, USA¹

Abstract

In this note we present a model that can produce a mass shift in a resonance due to interference between a scattering amplitude and that amplitude having rescattering through the resonance.

1 Starting point

For the first part of this story we will define what a scattering cross section is. We will only consider in this work elastic scattering of pions. Two pions can scatter at a certain energy which we will call $M_{\pi\pi}$. The differential cross section σ at a given $M_{\pi\pi}$ is

$$\frac{d\sigma}{d\phi d\theta} = \frac{1}{K^2} \left| \sum_{\ell} (2\ell + 1) T_{\ell} P_{\ell}(\cos(\theta)) \right|^2 \quad (1)$$

where ϕ and θ are the azimuthal and scattering angles, respectively. T_{ℓ} is a complex scattering amplitude and ℓ is the angular momentum. P_{ℓ} is the Legendre polynomial, which is a function of $\cos(\theta)$. K is the flux factor equal to the pion momentum in the center of mass. For this note we only consider $\ell = 0$ and $\ell = 1$. The T_0 and T_1 elastic scattering amplitudes are complex amplitudes described by one real number which is in units of angles. The form of the amplitude is

$$T_{\ell} = e^{i\delta_{\ell}} \sin(\delta_{\ell}) \quad (2)$$

We note that δ depends on the value of ℓ and $M_{\pi\pi}$. We will use the $\ell = 0$ and $\ell = 1$ δ 's given in [1].

¹This research was supported by the U.S. Department of Energy under Contract No. DE-AC02-98CH10886

2 Defining A

Let us consider two pions scattering in the final state of the heavy ion collision. The scattering will be either $\ell = 0$ or $\ell = 1$ partial waves. The $M_{\pi\pi}$ of the scattering di-pion system will depend on the probability of the phase space of the overlapping pions. The pions emerge from a close encounter in a defined quantum state with a random phase. We will call this amplitude A and note that the absolute value squared of the amplitude is proportional to the phase space overlap. The emerging pions can re-interact or re-scatter through the quantum state of the pions, which is a partial wave or a phase shift. We have amplitude A plus A times the re-scattering of pions through the phase shift consistent quantum state of A . The correct unitary way to describe this process is given by [2] equation (4.5)

$$T = \frac{V_1 U_1}{D_1} + \frac{\left(V_2 + \frac{D_{12} V_1}{D_1}\right) \left(U_2 + \frac{D_{12} U_1}{D_1}\right)}{D_2 - \frac{D_{12}^2}{D_1}} \quad (3)$$

In the above equation we have two terms, 1 and 2. The first term denoted by 1 is the $\pi\pi$ scattering through p-wave which will become the amplitude A mentioned above, where V is the incoming and U is the outgoing $\pi\pi$ system. The second term denoted by 2 is the direct production of the ρ meson with V being the production, the propagation being D and the decay being U . We see that there are terms D_{12} which involves a loop of pions between scattering pions and the formation of a ρ by the pions.

3 Final equation

The complete derivation is in the appendix. From the appendix we get two terms, one being the direct production of the ρ or $\pi\pi$ p-wave phase shift and the second being the ρ from re-scattering. The final equation number 21 has two important factors, one is two-body phase space and the other is a coefficient α . This coefficient is related to the real part of the $\pi\pi$ rescattering loop and is given by equation 14. When the pions rescatter or interact at a close distance or a point α has its maximum value of one. While if the pions rescatter or interact at a distance determined by the diffractive limit the value of α is zero.

In the equation 21 $|T|^2$ is the cross section for ρ production, where D is the direct production amplitude and A is the amplitude introduced above

for the re-scattering pions into the ρ meson. δ is the $\pi\pi$ phase shift [1]. q is the $\pi\pi$ center of mass. At a given p_T and y bin, D will be a constant as a function of $M_{\pi\pi}$. The α factor is one for rescattering coming from a point source, but goes to zero when rescattering is diffractive. The dependence of A is calculated by the phase space overlap of di-pions added as four vectors and corrected for proper time, with the sum having the correct p_T and y for a given $M_{\pi\pi}$.

Finally we must use the correct two body phase space. For a two body system of pions, phase space goes to a constant as $M_{\pi\pi}$ goes to infinity. Let us choose this constant to be unity. Phase space which is denoted by PS is equal to

$$PS = \frac{2qB_\ell(q/q_0)}{M_{\pi\pi}} \quad (4)$$

where B_ℓ is a Blatt-Weisskopf-barrier factor[3] for ℓ angular momentum quantum number. The q_0 is the momentum related to the range of interaction of the $\pi\pi$ scattering. 1 fm is the usual interaction distance which implies that q_0 is 200 MeV/c. For the ρ meson $\ell=1$ and $B_1 = \frac{(q/q_0)^2}{(1+(q/q_0)^2)}$.

The phase space factor PS as a function of q near the $\pi\pi$ threshold is given by $q^{2\ell+1}$. Thus in the appendix we use $q^{2\ell+1}$ for the factor PS except for equation 21 which is the final equation.

A Appendix

Starting with equation (4.5) from [2]

$$T = \frac{V_1 U'_1}{D_1} + \frac{\left(V_2 + \frac{D_{12} V_1}{D_1}\right) \left(U'_2 + \frac{D_{12} U'_1}{D_1}\right)}{D_2 - \frac{D_{12}^2}{D_1}} \quad (5)$$

In order to have the correct threshold kinematics, we define

$$U'_1 = U_1 \sqrt{q^{2\ell+1}} \quad (6)$$

$$U'_2 = U_2 \sqrt{q^{2\ell+1}} \quad (7)$$

where q is the $\pi\pi$ center of mass momentum and ℓ is the value of the angular momentum. The amplitude A of the text is given by

$$\frac{V_1 U_1}{D_1} = A \quad (8)$$

Thus we have

$$\frac{V_1 U'_1}{D_1} = A \sqrt{q^{2\ell+1}} \quad (9)$$

The phase shift for the ℓ^{th} partial wave will be given by δ_ℓ , where

$$\frac{U'_2 U'_2}{D_2} = e^{i\delta_\ell} \sin(\delta_\ell) \quad (10)$$

The above equality is true if the D_1 mode plays no role in the $\pi\pi$ scattering in the ℓ^{th} partial wave. But in the initial state there is a large production of D_1 . The U 's are the basic coupling of the D 's to the $\pi\pi$ system. In order to decouple D_1 from the $\pi\pi$ system U_1 must go to zero. We can maintain a finite production of D_1 if we define

$$V_1 = \frac{1}{U_1} \quad (11)$$

Thus the first term in the equation becomes

$$\frac{V_1 U'_1}{D_1} = \frac{\frac{1}{U_1} U_1 \sqrt{q^{2\ell+1}}}{D_1} = \frac{\sqrt{q^{2\ell+1}}}{D_1} = A \sqrt{q^{2\ell+1}} \quad (12)$$

The form of D_{12} is given by

$$D_{12} = \alpha U_1 U_2 + i q^{2\ell+1} U_1 U_2 \quad (13)$$

D_{12} is the real and imaginary part of the two pion loop from state 1 to state 2. The U 's are the $\pi\pi$ couplings and the imaginary part goes to zero at the $\pi\pi$ threshold. The α factor is one for rescattering coming from a point source, but goes to zero when rescattering is diffractive. A simple form for α is given by

$$\alpha = \left(1.0 - \frac{r^2}{r_0^2}\right) \quad (14)$$

where r is the radius of rescattering in fermis and r_0 is 1.0 fermi or the limiting range of the strong interaction. The second term of the first equation is

$$\frac{\left(V_2 + \frac{D_{12} V_1}{D_1}\right) \left(U'_2 + \frac{D_{12} U'_1}{D_1}\right)}{D_2 - \frac{D_{12}^2}{D_1}} \quad (15)$$

Rewriting

$$\frac{\left(V_2 + \frac{\alpha U_1 U_2 V_1}{D_1} + i q^{2\ell+1} \frac{U_1 U_2 V_1}{D_1}\right) \left(U'_2 + \frac{D_{12} U'_1}{D_1}\right)}{D_2 - \frac{D_{12}^2}{D_1}} \quad (16)$$

Let us make substitutions

$$V_1 = \frac{1}{U_1}, U_2 = \frac{U'_2}{\sqrt{q^{2\ell+1}}}, \frac{1}{D_1} = A, D_{12} = 0 \quad (17)$$

The second term becomes

$$\frac{\left(V_2 + \frac{A \alpha U'_2}{\sqrt{q^{2\ell+1}}} + i \sqrt{q^{2\ell+1}} A U'_2\right) U'_2}{D_2} \quad (18)$$

The first term is

$$\frac{V_1 U'_1}{D} = A \sqrt{q^{2\ell+1}} \quad (19)$$

Adding the first and the second terms and substituting the phase shift,

$$T = \frac{V_2}{U_2} \frac{e^{i\delta_\ell} \sin(\delta_\ell)}{\sqrt{q^{2\ell+1}}} + A \left(\frac{e^{i\delta_\ell} \alpha \sin(\delta_\ell)}{\sqrt{q^{2\ell+1}}} + \sqrt{q^{2\ell+1}} e^{i\delta_\ell} \cos(\delta_\ell) \right) \quad (20)$$

The term with the factor $\frac{V_2}{U_2}$ is the direct production of the di-pion system. We shall call this amplitude D . The re-scattered amplitude is A and is modified by the di-pion phase shift. These two amplitudes have some random phase and are not coherent. Thus the cross section is

$$|T|^2 = |D|^2 \frac{\sin^2(\delta_\ell)}{PS} + \frac{|A|^2}{PS} |\alpha \sin(\delta_\ell) + PS \cos(\delta_\ell)|^2 \quad (21)$$

References

- [1] G. Grayner *et. al.*, Nucl. Phys. B 75 (1974) 189.
- [2] R. Aaron and R. S. Longacre, Phys. Rev. D 24 (1981) 1207.
- [3] F. von Hippel and C. Quigg, Phys. Rev. 5 (1972) 624.